

Pathogenesis and Pathophysiology of the Cardiometabolic Syndrome

Erik P. Kirk, PhD; Samuel Klein, MD

The cardiometabolic syndrome represents a cluster of metabolic abnormalities that are risk factors for cardiovascular disease. The mechanism(s) responsible for developing the cardiometabolic syndrome is not known, but it is likely that multi-organ insulin resistance, which is a common feature of the cardiometabolic syndrome, is involved. Insulin resistance is an important risk factor for type 2 diabetes and can cause vasoconstriction and renal sodium reabsorption, leading to increased blood pressure. Alterations in adipose tissue fatty acid and adipokine metabolism are involved in the pathogenesis of insulin resistance. Excessive rates of fatty acid release into the bloodstream can impair the ability of insulin to stimulate muscle glucose uptake and suppress hepatic glucose production. Noninfectious systemic inflammation associated with adipocyte and adipose tissue macrophage cytokine production can also cause insulin resistance. In addition, increased free fatty acid delivery to the liver can stimulate hepatic very low-density lipoprotein triglyceride production, leading to dyslipidemia. J Clin Hypertens (Greenwich). 2009;11:761–765. ©2009 Wiley Periodicals, Inc.

From the Center for Human Nutrition, Washington University School of Medicine, St Louis, MO
Address for correspondence:
Samuel Klein, MD, Washington University School of Medicine, 660 South Euclid Avenue, Box 8031, St Louis, MO 63110-1093
E-mail: sklein@wustl.edu
Manuscript received June 19, 2008; revised October 10, 2008; accepted December 2, 2008

The cardiometabolic syndrome represents a constellation of metabolic abnormalities that are risk factors for cardiovascular disease. The risk of coronary heart disease, myocardial infarction, and stroke is much higher in persons who have the cardiometabolic syndrome than in those without the syndrome.¹ No universally accepted definition of the cardiometabolic syndrome has been established, and at least 5 independent groups have proposed clinical criteria for establishing its diagnosis.² The most widely used clinical criteria for diagnosing the cardiometabolic syndrome are those proposed by the World Health Organization³ and the National Cholesterol Education Program Adult Treatment Panel III (NCEP ATP III)⁴ (Table). The common characteristics of the cardiometabolic syndrome among all groups include abdominal obesity (high body mass index and/or large waist circumference), insulin-resistant glucose metabolism (hyperinsulinemia, impaired fasting glucose, impaired glucose tolerance, type 2 diabetes), dyslipidemia (high serum triglyceride and low serum high-density lipoprotein cholesterol concentrations), and increased blood pressure.

PREVALENCE

The cardiometabolic syndrome has become a major public health problem in the United States and many other countries worldwide because of its increasing prevalence. Data from the third National Health and Nutrition Examination Survey (NHANES) (1988–1994) found that the age-adjusted prevalence of the cardiometabolic syndrome, defined by using the ATP III criteria, was 24% in the adult US population.⁴ The prevalence of the cardiometabolic syndrome increases linearly

doi: 10.1111/j.1559-4572.2009.00054.x



Table. Clinical Identification of the Cardiometabolic Syndrome Based on Criteria From the NCEP ATP III or the WHO

	NCEP ATP III ^a	WHO ^b
Fasting blood glucose	≥100 mg/dL	IFG/IGT/T2DM
Abdominal obesity		
Men	>102 cm WC	>0.90 WHR (or BMI ≥30 kg/m ²)
Women	>88 cm WC	>0.85 WHR (or BMI ≥30 kg/m ²)
Triglycerides	≥150 mg/dL	≥1.7 mmol L ⁻¹
HDL Cholesterol		
Men	<40 mg/dL	<0.9 mmol L ⁻¹
Women	<50 mg/dL	<1.0 mmol L ⁻¹
Blood pressure	≥130/85 mm Hg	≥140/90 mm Hg
Microalbuminuria	–	Yes

Abbreviations: BMI, body mass index; HDL, high-density lipoprotein; IFG, impaired fasting glucose; IGT, impaired glucose tolerance; NCEP ATP III, National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults Adult Treatment Panel III; T2DM, type 2 diabetes mellitus; WC, waist circumference; WHO, World Health Organization; WHR, waist-to-hip circumference ratio. ^aThree or more criteria. ^bIFG/IGT/T2DM plus ≥2 criteria.




Figure 1. Physiologic interrelationships between fatty acid metabolism, insulin resistance, and features of the cardiometabolic syndrome. CETP indicates cholesterol ester transfer protein; VLDL, very low-density lipoprotein triglyceride; HDL, high-density lipoprotein; TG, triglyceride; FFA, free fatty acid.

with age from approximately 7% in those who are 20 to 29 years old to approximately 45% in those who are 60 years and older. Moreover, the latest NHANES data found that the prevalence of the cardiometabolic syndrome is increasing in both men and women in all age groups.⁴

PATHOPHYSIOLOGY

Fatty Acid Metabolism

The cardiometabolic syndrome is also known as the *insulin resistance syndrome* because it has been hypothesized that insulin resistance is the major mechanism responsible for the metabolic abnormalities of the syndrome.⁵ Alterations in free fatty acid metabolism are likely a major factor involved in the pathogenesis of hyperglycemia and dyslipidemia associated with the cardiometabolic syndrome (Figure 1). Excessive release of free fatty acids from adipose tissue into plasma and increased plasma

free fatty acid concentration can impair the ability of insulin to stimulate muscle glucose uptake⁶ and suppress hepatic glucose production.⁷ In addition, increased free fatty acid delivery to the liver can increase hepatic very low-density lipoprotein triglyceride production^{8,9} and plasma triglyceride concentration.¹⁰ An increase in plasma triglycerides increases the transfer of triglycerides from very low-density lipoprotein to high-density lipoprotein, which leads to increased high-density lipoprotein clearance and decreased plasma high-density lipoprotein concentration.¹¹

Insulin, which inhibits lipolysis, is the major physiologic regulator of basal adipose tissue lipolytic activity.^{12,13} Lipolysis of adipose tissue triglycerides is the major source of plasma free fatty acids.¹⁴ Therefore, insulin resistance in adipose tissue stimulates an increase in lipolytic rate and free fatty acid release into the bloodstream. The typical increase in plasma insulin concentrations associated with obesity does not completely compensate for adipose tissue insulin resistance, so insulin-resistant obese persons have high basal lipolytic rates and plasma free fatty acid concentrations.¹³

In skeletal muscle, the cellular mechanism responsible for free fatty acid-induced insulin resistance involves alterations in intracellular insulin signaling and impaired insulin-mediated glucose uptake^{15,16} (Figure 2). An acute increase in plasma free fatty acid concentrations from approximately 400 $\mu\text{mol L}^{-1}$ (normal basal concentration) to approximately 800 $\mu\text{mol L}^{-1}$ (concentration during short-term fasting) causes a marked increase in intramyocellular fatty acid metabolites, including long-chain fatty acyl-CoA and diacylglycerol.^{15,17,18} These

metabolites are potent allosteric activators of protein kinase C, a serine/threonine kinase that phosphorylates serine/threonine sites of insulin receptor substrate-1, thereby inhibiting insulin's ability to activate phosphoinositide 3-kinase activity¹⁹⁻²¹ and decreasing downstream events, including translocation of glucose transporter 4 from the cytoplasm to the cell membrane needed for glucose transport.

Other factors related to intracellular fatty acid metabolism can also contribute to insulin resistance (Figure 2). Defective skeletal muscle mitochondrial function has been identified in persons who have insulin resistance and are at increased risk for type 2 diabetes.²² Impaired mitochondrial fatty acid oxidation can contribute to impaired insulin action by increasing the intracellular accumulation of fatty acids. In addition, excessive intracellular fatty acids can increase the production of reactive oxygen species, which leads to activation of the proinflammatory nuclear factor kappa B pathway,^{17,23} thereby increasing insulin resistance.

The cellular events responsible for fatty acid-induced insulin resistance in the liver have not been as carefully evaluated as in skeletal muscle. Increased delivery of free fatty acids to the liver and possibly increased release of fatty acids from lipolysis of intrahepatic triglycerides stimulate hepatic glucose production.¹⁹⁻²¹ Free fatty acid-induced insulin resistance in the liver is associated with activation of protein kinase C.²⁴

Abdominal Adipose Tissue

Excess abdominal fat mass, particularly visceral (intrahepatic) fat, is associated with insulin resistance.^{6,7,25,26} However, it is not known whether visceral fat causes or is simply associated with insulin resistance. Visceral fat represents a small component of total body fat mass. Visceral fat accounts for about 10% of total body fat mass in lean men and about 15% of total body fat mass in obese men.²⁶ Nonetheless, it has been hypothesized that fatty acids released during lipolysis of visceral adipose tissue are an important cause of insulin resistance because these fatty acids enter the portal vein and are delivered directly to the liver.²⁷ Data from studies that used isotope tracers to assess visceral fat metabolism in vivo in obese persons found that approximately 20% of free fatty acids delivered to the liver and approximately 15% of free fatty acids delivered to skeletal muscle are derived from lipolysis of visceral fat.²⁸ Therefore, visceral fat might contribute to hepatic insulin resistance, but it is unlikely that visceral fat is responsible for insulin resistance in skeletal muscle.




Figure 2. Potential cellular mechanisms for fatty acid-induced insulin resistance. IRS indicates insulin receptor substrate; PI 3 kinase, phosphoinositide 3-kinase; PKC, protein kinase C; TG, triglyceride; ROS, reactive oxygen species; IKK- β , I-kappa B kinase β ; NF κ B, nuclear factor kappa B. Adapted from Shulman.¹⁶

Ectopic Fat

Ectopic accumulation of fat in liver and muscle cells is associated with insulin resistance in those tissues.^{20,29} Increased intrahepatic fat content is associated with hepatic insulin resistance in the liver and impaired insulin-mediated suppression of hepatic glucose production,²⁰ and increased intramyocellular fat content is associated with skeletal muscle insulin resistance and impaired insulin-mediated glucose disposal.²⁹

Adipose Tissue Secretory Proteins

Adipose tissue produces several inflammatory cytokines (adipokines), which can induce insulin resistance, and adiponectin, which increases insulin sensitivity.^{23,30} For example, tumor necrosis factor α suppresses insulin signaling,³¹ interleukin-6 increases inflammation directly or by stimulating hepatic C-reactive protein production,³² macrophage chemoattractant protein 1 is a potent chemoattractant for macrophages,³³ and interleukin-8 activates neutrophil granulocytes and is chemotactic for all known migratory immune cells.³⁴ Adiponectin increases insulin sensitivity in the liver, decreases hepatic glucose production,³⁵ and increases skeletal muscle glucose and fatty acid oxidation.³⁶

Increased Blood Pressure

The relationship between insulin resistance and hypertension is well established.³⁷ Fatty acids themselves can cause vasoconstriction.³⁸ Additionally, insulin resistance can increase blood pressure because insulin is a vasodilator,³⁹ and hyperinsulinemia increases renal sodium reabsorption.⁴⁰ Persons who are insulin-resistant tend to lose the vasodilatory

effect of insulin⁴¹ but preserve the renal effect on sodium reabsorption,⁴⁰ and sodium reabsorption is increased in persons with the cardiometabolic syndrome.⁴²

CONCLUSIONS

The cardiometabolic syndrome includes a cluster of conditions including abdominal obesity, insulin-resistant glucose metabolism, dyslipidemia, and increased blood pressure. Alterations in fatty acid metabolism (eg, excessive fatty acid release into plasma) likely contribute to these metabolic abnormalities. Increased free fatty acids can (1) impair insulin action in skeletal muscle and liver, leading to increased blood glucose concentration; (2) stimulate hepatic very low-density lipoprotein triglyceride production, leading to increased serum triglyceride and decreased high-density lipoprotein concentrations; and (3) stimulate vasoconstriction and increase sodium reabsorption, possibly leading to increased blood pressure.

REFERENCES

- 1 Isomaa B, Almgren P, Tuomi T, et al. Cardiovascular morbidity and mortality associated with the metabolic syndrome. *Diabetes Care*. 2001;24:683–689.
- 2 Grundy SM, Cleeman JI, Daniels SR, et al. Diagnosis and management of the metabolic syndrome. An American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. Executive summary. *Cardiol Rev*. 2005;13:322–327.
- 3 *Obesity: Preventing and Managing the Global Epidemic: Report of a WHO Consultation*. Geneva, Switzerland: World Health Organization; 2000.
- 4 Expert panel on detection, evaluation, and treatment of high blood cholesterol in adults. Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). *JAMA*. 2001;285:2486–2497.
- 5 Reaven GM. Banting lecture 1988. Role of insulin resistance in human disease. *Diabetes*. 1988;37:1595–1607.
- 6 Abate N, Garg A, Peshock RM, et al. Relationship of generalized and regional adiposity to insulin sensitivity in men with NIDDM. *Diabetes*. 1996;45:1684–1693.
- 7 Pouliot MC, Despres JP, Nadeau A, et al. Visceral obesity in men. Associations with glucose tolerance, plasma insulin, and lipoprotein levels. *Diabetes*. 1992;41:826–834.
- 8 Lewis GF, Uffelman KD, Szeto LW, et al. Interaction between free fatty acids and insulin in the acute control of very low density lipoprotein production in humans. *J Clin Invest*. 1995;95:158–166.
- 9 Zhang YL, Hernandez-Ono A, Ko C, et al. Regulation of hepatic apolipoprotein B-lipoprotein assembly and secretion by the availability of fatty acids. I. Differential response to the delivery of fatty acids via albumin or remnant-like emulsion particles. *J Biol Chem*. 2004;279:19362–19374.
- 10 Mittendorfer B, Patterson BW, Klein S. Effect of sex and obesity on basal VLDL-triacylglycerol kinetics. *Am J Clin Nutr*. 2003;77:573–579.
- 11 Hopkins GJ, Barter PJ. Role of triglyceride-rich lipoproteins and hepatic lipase in determining the particle size and composition of high density lipoproteins. *J Lipid Res*. 1986;27:1265–1277.
- 12 Klein S, Peters EJ, Holland OB, et al. Effect of short- and long-term beta-adrenergic blockade on lipolysis during fasting in humans. *Am J Physiol Endocrinol Metab*. 1989;257:E65–E73.
- 13 Horowitz JF, Klein S. Whole body and abdominal lipolytic sensitivity to epinephrine is suppressed in upper body obese women. *Am J Physiol Endocrinol Metab*. 2000;278:E1144–E1152.
- 14 Mittendorfer B, Liem O, Patterson BW, et al. What does the measurement of whole-body fatty acid rate of appearance in plasma by using a fatty acid tracer really mean? *Diabetes*. 2003;52:1641–1648.
- 15 Boden G, Lebed B, Schatz M, et al. Effects of acute changes of plasma free fatty acids on intramyocellular fat content and insulin resistance in healthy subjects. *Diabetes*. 2001;50:1612–1617.
- 16 Shulman GI. Cellular mechanisms of insulin resistance. *J Clin Invest*. 2000;106:171–176.
- 17 Itani SI, Ruderman NB, Schmieder F, et al. Lipid-induced insulin resistance in human muscle is associated with changes in diacylglycerol, protein kinase C, and IkkappaB-alpha. *Diabetes*. 2002;51:2005–2011.
- 18 Yu C, Chen Y, Cline GW, et al. Mechanism by which fatty acids inhibit insulin activation of insulin receptor substrate-1 (IRS-1)-associated phosphatidylinositol 3-kinase activity in muscle. *J Biol Chem*. 2002;277:50230–50236.
- 19 Kelley DE, Moka M, Simoneau JA, et al. Interaction between glucose and free fatty acid metabolism in human skeletal muscle. *J Clin Invest*. 1993;92:91–98.
- 20 Seppala-Lindroos A, Vehkavaara S, Hakkinen AM, et al. Fat accumulation in the liver is associated with defects in insulin suppression of glucose production and serum free fatty acids independent of obesity in normal men. 10.1210/jc.87.7.3023. *J Clin Endocrinol Metab*. 2002;87:3023–3028.
- 21 Ferrannini E, Barrett EJ, Bevilacqua S, et al. Effect of fatty acids on glucose production and utilization in man. *J Clin Invest*. 1983;72:1737–1747.
- 22 Petersen KF, Dufour S, Befroy D, et al. Impaired mitochondrial activity in the insulin-resistant offspring of patients with type 2 diabetes. *N Engl J Med*. 2004;350:664–671.
- 23 Steinberg HO, Tarshoby M, Monestel R, et al. Elevated circulating free fatty acid levels impair endothelium-dependent vasodilation. *J Clin Invest*. 1997;100:1230–1239.
- 24 Lam TK, Yoshii H, Haber CA, et al. Free fatty acid-induced hepatic insulin resistance: a potential role for protein kinase C-delta. *Am J Physiol Endocrinol Metab*. 2002;283:E682–E691.
- 25 Ross R, Aru J, Freeman J, et al. Abdominal adiposity and insulin resistance in obese men. *Am J Physiol Endocrinol Metab*. 2002;282:E657–E663.
- 26 Abate N, Garg A, Peshock RM, et al. Relationships of generalized and regional adiposity to insulin sensitivity in men. *J Clin Invest*. 1995;96:88–98.
- 27 Nielsen S, Guo Z, Johnson CM, et al. Splanchnic lipolysis in human obesity. *J Clin Invest*. 2004;113:1582–1588.
- 28 Havel RJ, Kane JP, Balasse EO, et al. Splanchnic metabolism of free fatty acids and production of triglycerides of very low density lipoproteins in normotriglyceridemic and hypertriglyceridemic humans. *J Clin Invest*. 1970;49:2017–2035.
- 29 Krssak M, Falk Petersen K, Dresner A, et al. Intramyocellular lipid concentrations are correlated with insulin sensitivity in humans: a 1H NMR spectroscopy study. *Diabetologia*. 1999;42:113–116.

- 30 Perseghin G, Petersen K, Shulman GI. Cellular mechanism of insulin resistance: potential links with inflammation. *Int J Obes Relat Metab Disord*. 2003;27(suppl 3):S6–S11.
- 31 Crespo J, Cayon A, Fernandez-Gil P, et al. Gene expression of tumor necrosis factor alpha and TNF-receptors, p55 and p75, in nonalcoholic steatohepatitis patients. *Hepatology*. 2001;34:1158–1163.
- 32 Park SH, Kim BI, Yun JW, et al. Insulin resistance and C-reactive protein as independent risk factors for non-alcoholic fatty liver disease in non-obese Asian men. *J Gastroenterol Hepatol*. 2004;19:694–698.
- 33 Christiansen T, Richelsen B, Bruun JM. Monocyte chemoattractant protein-1 is produced in isolated adipocytes, associated with adiposity and reduced after weight loss in morbid obese subjects. *Int J Obes (Lond)*. 2005;29:146–150.
- 34 Baggiolini M, Loetscher P, Moser B. Interleukin-8 and the chemokine family. *Int J Immunopharmacol*. 1995;17:103–108.
- 35 Berg AH, Combs TP, Du X, et al. The adipocyte-secreted protein Acrp30 enhances hepatic insulin action. *Nat Med*. 2001;7:947–953.
- 36 Yamauchi T, Kamon J, Minokoshi Y, et al. Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase. *Nat Med*. 2002;8:1288–1295.
- 37 Ferrannini E, Buzzigoli G, Bonadonna R, et al. Insulin resistance in essential hypertension. *N Engl J Med*. 1987;317:350–357.
- 38 Tripathy D, Mohanty P, Dhindsa S, et al. Elevation of free fatty acids induces inflammation and impairs vascular reactivity in healthy subjects. *Diabetes*. 2003;52:2882–2887.
- 39 Steinberg HO, Brechtel G, Johnson A, et al. Insulin-mediated skeletal muscle vasodilation is nitric oxide dependent. A novel action of insulin to increase nitric oxide release. *J Clin Invest*. 1994;94:1172–1179.
- 40 DeFronzo RA, Cooke CR, Andres R, et al. The effect of insulin on renal handling of sodium, potassium, calcium, and phosphate in man. *J Clin Invest*. 1975;55:845–855.
- 41 Tooke JE, Hannemann MM. Adverse endothelial function and the insulin resistance syndrome. *J Intern Med*. 2000;247:425–431.
- 42 Barbato A, Cappuccio FP, Folkard EJ, et al. Metabolic syndrome and renal sodium handling in three ethnic groups living in England. *Diabetologia*. 2004;47:40–46.